

05/26/99



Jc571 U.S. PTO

A

Jc588 U.S. PTO  
09/320637

05/26/99

**UTILITY  
PATENT APPLICATION  
TRANSMITTAL**

(Only for new nonprovisional applications under 37 CFR 1.53(b))

Attorney Docket No. 3339-239A

First Inventor or Application Identifier: Joël Sternheimer

Title of Invention: METHOD FOR THE REGULATION OF  
PROTEIN BIOSYNTHESIS

Express Mail Label No. EL110131547US

ADDRESS TO: ASSISTANT COMMISSIONER FOR PATENTS  
BOX PATENT APPLICATION  
WASHINGTON, DC 20231

Transmitted herewith for filing in the United States Patent Office is a patent application for:

Inventors: Joël Sternheimer

1. ☒ The Filing Fee has been calculated as shown below:  
(Submit an original, and a duplicate for fee processing)

	No. Filed	No. Extra	Small Entity		Large Entity	
			Rate	Fee 1	Rate	Fee 0
BASIC FEE				\$380		\$0
TOTAL CLAIMS:	12 - 20 =	0		X 9 = \$0		x 18 = \$0
INDEP CLAIMS:	1 - 3 =	0		X 39 = \$0		x 78 = \$0
<input type="checkbox"/> MULTIPLE DEPENDENT CLAIMS PRESENTED				+130 = \$		+260 = \$
*If the difference in Column 1 is less than zero, enter "0" in Column 2.				TOTAL \$ 380		TOTAL \$

The Commissioner is hereby authorized to credit overpayments or charge the following fees to Deposit Account No. 16-0605.

- a. ☒ Fees required under 37 CFR 1.16 (National filing fees).  
b. ☒ Fees required under 37 CFR 1.17 (National application processing fees).  
☒ A check in the amount of \$ 380.00 is enclosed.  
☐ The above filing fee will be paid along with Applicant(s) Response to the Notice to File Missing Parts.
2. ☒ Specification; Total Pages 34
3. ☒ 3 Sheets of Formal Drawing(s) (35 USC 113)

4. ☒ Declaration and Power of Attorney; [Total Pages 3]
- a. ☒ Newly executed (original or copy)
- b. ☐ Copy from a prior application (37 CFR 1.63(d))  
(for continuation/divisional with Box 16 completed)
- i. ☐ DELETION OF INVENTOR(S) Signed statement attached deleting inventor(s) named in the prior application, see 37 CFR 1.63(d)(2) & 1.33(b).
5. ☐ Microfiche Computer Program (Appendix)
6. ☐ Nucleotide and/or Amino Acid Sequence Submission (if applicable, all necessary)
- a. ☐ Computer Readable Copy
- b. ☐ Paper Copy (identical to computer copy)
- c. ☐ Statement verifying identity of above copies

#### ACCOMPANYING APPLICATION PARTS

7. ☐ Assignment Papers (cover sheet & document(s) (including \$40.00 fee)
8. ☐ 37 CFR 3.73(b) Statement (when there is an assignee); ☐ Power of Attorney
9. ☐ English Translation Document (if applicable)
10. ☒ Information Disclosure Statement (IDS)/PTO-1449; 0 Copies of IDS Citations
11. ☒ Preliminary Amendment
12. ☒ Return Receipt Postcard (MPEP 503) (Should be specifically itemized)
13. ☒ Small Entity Statement(s)  
☐ Statement filed in prior application; status still proper and desired.
14. ☐ Certified Copy of Priority Document(s) (if foreign priority is claimed)  
Foreign Priority is
15. ☒ Other: Rule 132 Declaration Of Joel Sternheimer

16. **If a CONTINUING APPLICATION, check appropriate box and supply the requisite information below and in a preliminary amendment:**

- ☐ Continuation ☐ Divisional ☒ Continuation in Part (CIP)  
of prior application No: 08/347,353; Filed December 1, 1994

Prior Application Information: Examiner J. Martinell Group/Art Unit: 1804

For CONTINUATION or DIVISIONAL APPS only: The entire disclosure of the prior application, from which an oath or declaration is supplied under Box 4b, is considered a part of the disclosure of the accompanying continuation or divisional application and is hereby incorporated by reference. The incorporation can only be relied upon when a portion has been inadvertently omitted from the submitted application parts.

17. **CORRESPONDENCE ADDRESS**

Customer Number or Bar Code Label **000826**

Attention Of: Ernest B. Lipscomb, III

Signature:

*Ernest B. Lipscomb III*

Attorney of Record: Ernest B. Lipscomb, III

Attorney Registration No. 24,733

Tel Charlotte Office (704) 331-6000

Fax Charlotte Office (704) 334-2014

**ALSTON & BIRD LLP**

P.O. Drawer 34009

Charlotte NC 28234-4009

CLT01/4369142v1

"Express Mail" mailing label number EL110131547US

Date of Deposit: May 26, 1999

I hereby certify that this paper or fee is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 CFR 1.10 on the date indicated above and is addressed to Box Patent Application, Assistant Commissioner For Patents, Washington, DC 20231.

*Kim M. Ramsey*

Kim M. Ramsey

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Attorney's Docket No. 3339-239A

Applicant, Patentee, or Identifier: Joel Sternheimer  
Application No. or Patent No.: To Be Assigned  
Filed or Issued: Concurrently Herewith  
Title: METHOD FOR THE REGULATION OF  
PROTEIN BIOSYNTHESIS

STATEMENT CLAIMING SMALL ENTITY STATUS  
(37 C.F.R. § 1.9(f) and 1.27(b))--INDEPENDENT INVENTOR

As a below named inventor, I hereby declare that I qualify as an independent inventor as defined in 37 C.F.R. § 1.9(c) for purposes of paying reduced fees under section 41(a) and (b) of Title 35, United States Code, to the Patent and Trademark Office described in:

- ☒ the specification filed herewith with title as listed above.  
☐ the application identified above.  
☐ the patent identified above.

I have not assigned, granted, conveyed, or licensed, and am under no obligation under contract or law to assign, grant, convey, or license, any rights in the invention to any person who would not qualify as an independent inventor under 37 C.F.R. § 1.9(c) if that person had made the invention, or to any concern which would not qualify as a small business concern under 37 C.F.R. § 1.9(d) or nonprofit organization under 37 C.F.R. § 1.9(e).

Each person, concern, or organization to which I have assigned, granted, conveyed, or licensed or am under an obligation under contract or law to assign, grant, convey, or license any rights in the invention is listed below:

- ☒ No such person, concern, or organization exists.  
☐ Each such person, concern, or organization is listed below.

FULL NAME: \_\_\_\_\_

ADDRESS: \_\_\_\_\_

☐ Individual

☐ Small Business

☐ Nonprofit Organization

FULL NAME: \_\_\_\_\_

ADDRESS: \_\_\_\_\_

☐ Individual

☐ Small Business

☐ Nonprofit Organization

Separate statements are required from each named person, concern, or organization having rights to the invention stating their status as small entities. (37 C.F.R. § 1.27)

I acknowledge the duty to file, in this application or patent, notification of any change in status resulting in loss of entitlement to small entity status prior to paying, or at the time of paying, the earliest of the issue fee or any maintenance fee due after the date on which status as a small entity is no longer appropriate. (37 C.F.R. § 1.28(b))

Joel Sternheimer  
NAME OF INVENTOR

Joel Sternheimer JSN  
(Signature of Inventor)

12/5/95  
Date

CLT01/4358044v1

Attorney's Docket No. 3339-239A

**PATENT**

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re: Joël Sternheimer  
Appl. No.: To Be Assigned  
Filed: Concurrently Herewith  
For: METHOD FOR THE REGULATION OF  
PROTEIN BIOSYNTHESIS

May 26, 1999

Assistant Commissioner for Patents  
Washington, DC 20231

**PRELIMINARY AMENDMENT**

Dear Sir:

Consideration of this Preliminary Amendment is respectfully requested. The Applicant has filed a continuation-in-part application in an attempt to more clearly set forth the subject invention in a manner that can be readily understood.

One of the issues in the parent application involved whether or not the claimed invention had utility within the meaning of 35 U.S.C. § 101.


The Applicant has submitted herewith a Rule 132 Declaration in which he has set forth a plethora of examples of practical utility (Annexes 3-6 and 9) of the claimed invention and testimonials (Annexes 7-8). Consideration of Mr. Sternheimer's Declaration filed under 37 C.F.R. § 1.132 is respectfully requested.

In re: Joël Sternheimer  
Appl. No.: To Be Assigned  
Filed: Concurrently Herewith  
Page 2

In the parent application a number of terms or phrases were rejected as being indefinite. The Applicant has dramatically revised the claims in a manner to eliminate or clarify both, if not all, of the objected to phrases in the claims of the parent application.

To the extent that the Examiner finds any indefiniteness or uncertainty in the claims, the Applicant has addressed each of these issues with clarifying the revisions to the specification.

Respectfully submitted,

  
Ernest B. Lipscomb, III  
Registration No. 24,7333

**ALSTON & BIRD LLP**

P.O. Drawer 34009

Charlotte, NC 28234

Tel Charlotte Office (704) 331-6000

Fax Charlotte Office (704) 334-2014

"Express Mail" mailing label number EL110131547US

Date of Deposit: May 26, 1999

I hereby certify that this paper or fee is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 CFR 1.10 on the date indicated above and is addressed to Box Patent Application, Assistant Commissioner For Patents, Washington, DC 20231.

  
Kim M. Ramsey

## METHOD FOR THE REGULATION OF PROTEIN BIOSYNTHESIS

### CROSS-REFERENCE TO RELATED APPLICATION

This application is a continuation-in-part of U.S.  
Patent Application, Serial No. 08/347,353 filed  
December 1, 1994.

### 5 BACKGROUND OF THE INVENTION

The present invention is directed to a method of  
regulating protein biosynthesis. More particularly, the  
invention is directed to a method for epigenetic  
regulation of *in situ* protein biosynthesis and its use in  
10 agronomy and health.

Demonstration of the musical properties of  
elementary particles suggests an important role for the  
scale at which the phenomena happen. (J. Sternheimer, C.  
R. Acad. Sc. Paris 297, 829, 1983). For example, it is  
15 known that the physical existence of quantum waves  
associated to particles propagate themselves not only in  
space-time, but also in that scale dimension, thus  
linking together successive levels of the organization of  
matter. (J. Sternheimer, *Colloque International "Louis de*  
20 *Broglie, Physicien et Penseur*", Ancienne Ecole  
Polytechnique, Paris, November 5-6, 1987). These waves  
allow an action of one scale onto the other, between  
phenomena that are similar enough to constitute, in a  
mathematically well-defined sense, harmonics of a common



fundamental tone. (See J. Sternheimer, *Ondes d'e'chelle* [scaling waves], I. Partie Physique; II. Partie Biologique. Filed at Academie des Sciences (Paris) 1992 under seal no. 17064).

5           The theoretical reasons for the existence of scaling waves makes them appear as a universal phenomenon whose function is at first to ensure coherence between the different scales of a quantum system, and that especially takes shape and can be described in the process of  
10 protein biosynthesis. The peptidic chain elongation effectively results from the sequential addition of amino acids that have been brought onto the ribosome by specific transfer RNAs (tRNAs). When an amino acid, initially in a free state, comes to affix itself to its  
15 tRNA, it is stabilized with respect to thermal agitation -- while keeping a relative autonomy because it is linked to the tRNA by only one degree of freedom -- for its de Broglie wavelength to reach the order of magnitude of its size. This stabilization gives the amino acid wave  
20 properties.

Interference between the scaling wave associated to the amino acid and those similarly produced by the other amino acids, results in a synchronization, after a very short period of time (which can be evaluated to be about  
25  $10^{-12.5}$  second), of the proper frequencies associated with these amino acids according to one and same musical scale, which more precisely depends upon the transfer RNA population. However, to within the approximation of the chromatic tempered scale, this scale appears universal  
30 due to the very peculiar distribution of amino acid masses which is already very close to it.

The scaling wave phenomenon appears in a more explicit way when the amino acid carried by its tRNA fixes itself onto the ribosome. It is at this moment that the stabilization with respect to thermal agitation becomes such that the wavelength of the amino acid outgrows its size by a full order of magnitude. The scaling wave which then emits interferes, at the scale of the protein in formation, with similar waves previously emitted by the other amino acids. This interference draws constraints of a musical type for the temporal succession of the proper frequencies associated to these waves, so that the scaling waves continue their itinerary and insure coherence and communication between different levels of the organism. For example, the succession of these waves minimizes the dissonance (harmonic distance) and the frequency gaps (represented by melodic distance) between successive amino acids. Additional properties imply the existence of periods of minimization of harmonic distances showing punctuations in the temporal succession of frequencies which other levels will complete with correlations all the more rich and marked that they themselves are more numerous to influence the protein synthesis. The result is the prediction that proteins possess, in the very succession of the proper quantum frequencies associated to the sequence of their amino acids, 'musical' properties all the more clear and elaborate that their biosynthesis is more sensitive to epigenetic factors in general. Conversely, it must be possible to act epigenetically, in a specific way for each protein onto that biosynthesis.

5 The observation of protein sequences confirms that  
all proteins possess musical properties in the sequence  
of their amino acids and these properties are all the  
more developed that those proteins are, in a general way,  
more epigenetically sensitive. (Data from M.O. Dayhoff,  
*Atlas of protein sequence and structure*, volume 5 and  
supplements, N.B.R.F. (Washington) 1972-78). In  
addition, the acoustic transposition of the series of  
proper frequencies corresponding to the production of  
10 scaling waves in phase with the elongation of a given  
protein, shows a stimulating action onto the biosynthesis  
of this protein *in vivo*, and in a correlative way it has  
an inhibiting action for scaling waves in phase  
opposition.

15 In the case of animals having a nervous system the  
sound wave is transformed into electromagnetic impulses  
of the same shape and frequency right from the starting  
point of the auditory nerve. These impulses, by virtue of  
the scale invariance of scaling wave equations applied to  
20 the photon (which generalize Maxwell's equations), have a  
direct action, by scale resonance, on their quantum  
transpositions. Because the squared quantum amplitudes  
are proportional to the number of proteins that are  
simultaneously synthesized, the resonance phenomenon  
25 results, in the case of scaling waves in phase, in an  
increase of the rate of synthesis, as well as a  
regulation of its rhythm, and in the case of scaling  
waves in phase opposition, in a reduction of this rate.  
(cf. P. Buser and M. Imbert, *Audition*, Hermann éditeur,  
30 Paris, 1987). Among plants, the sensitivity to sounds is

visible through interferometry and the scaling waves  
behave theoretically in a similar way.

The solution to the scaling wave equation, which  
effectively shows the existence of scaling waves having a  
range close to Avogadro number, anticipates similar  
properties for the scaling waves drawn from the spatial  
distribution of amino acids (whose de Broglie wavelength  
is then comparable to their size) inside the protein  
after it has been synthesized. The solution then  
provides a range approximating the square root of that  
number. The observation of their tertiary structures  
confirms the existence of harmonies within vibratory  
frequencies of amino acids spatially nearby inside  
proteins (and especially at their surface, as can be  
expected from their wavelength). An appreciable  
stabilization of the effects obtained with the use of the  
musical transpositions is then observed using colored  
transpositions of these spatially distributed  
frequencies.

The present invention is drawn from these  
observations.

#### SUMMARY OF THE INVENTION

The method of the invention comprises determining  
the musical notes associated with an amino acid sequence,  
the musical periods of the sequence, the lengths of the  
notes, and the tone quality of the notes through the  
retroaction of the amino acids and using that information  
to regulate the biosynthesis of the protein.

Stated in another way, the amino acids which build a  
protein emit a signal of quantum nature at a certain

frequency. Following the properties of this signal the frequency is transposed into a musical note in such way that playing back the melody of a protein will stimulate or inhibit its synthesis. This discovery has numerous applications since deduction of the amino acid sequence of a protein provides a sequence of notes composing the melody which will act on its synthesis inside an organism. Thus, by diffusing to a plant the music of a protein which plays an important role in flowering, more flowers are produced.

Stated more scientifically, the method of this invention uses the regulating action on the biosynthesis of proteins by scale resonance of transpositions into sound of temporal sequences of quantum vibrations associated with their elongation. This action may be an increase of the rate of synthesis or a reduction of this rate, depending upon whether the modulation of the vibration frequencies used is in phase with, or in phase opposition to the elongation. This is true for the quantum vibrations as well as for their transposition into sound. The result is further stabilized by the actions, again through scale resonance, of colored light transpositions of grouped quantum vibrations arising from the spatial conformation of proteins issued from this elongation.

This method applies in a specific way to every protein of known structure. Its use is all the more appropriate when the synthesis of this protein is even more dependent upon epigenetic factors, that is to say external to the DNA of the system to which it belongs, and especially in the present case, upon acoustic and

electromagnetic factors. In addition, the method uses the determination of metabolic agonisms and antagonisms of these proteins due to scale resonance phenomena naturally associated with their biosynthesis. The  
5 characterization of these proteins in their associated metabolic subsets is another feature of the present invention.

The identification of proteins designed to be regulated as part of a given application includes other  
10 criteria a correspondence between acoustic and electromagnetic phenomena or which effects can be observed on living beings and the transposed proteic sequences.

#### BRIEF DESCRIPTION OF THE INVENTION

15 Certain features and advantages will be evidence from the drawings when considered in conjunction with the accompanying drawing in which:

Fig. 1 shows the musical scale cytochrome oxidaze and cytochrome C;

20 Fig. 2 shows the cytochrome C humain region for amino-terminal and legends;

Fig. 3 shows Hystone IV and chalconesynthase; and

Fig. 4 shows "heat shock" HSP 27 Ethsp 70 and Troponinec.

#### 25 DETAILED DESCRIPTION OF THE INVENTION

The present invention will now be described more fully hereinafter with reference to the accompanying drawings, in which preferred embodiments of the invention are shown. This invention may, however, be embodied in

many different forms and should not be construed as limited to the embodiments set forth herein; rather, these embodiments are provided so that this disclosure will be thorough and complete, and will convey the scope of the invention to those skilled in the art.

There is provided a method of regulating protein synthesis *in situ*, using a musical sequence corresponding to the amino acid sequence of a protein through the decoding and transposition into sound of a temporal series of quantum vibrations associated with the elongation of the amino acid chain of the protein. The method of regulating protein synthesis *in situ* requires at least the following steps: the sequence of musical notes is determined; the period appearing in the molecule is determined; the period is rectified, if necessary; the rhythmic style is checked through the distribution of the bases of DNA; and the tone quality is determined.

Determining The Sequence Of Musical Notes. The sequent of music notes associated with the amino acid chain of a protein is determined by associating a musical note with each amino acid. More specifically, within the approximation of the tempered scale a universal code for the stimulation of protein synthesis is obtained. That code is:

Gly = low A; Ala = C; Ser = E; Pro Val, Thr, Cys = F; Leu, Ile, Asn, Asp = G; Gln, Lys, Glu, Met = A; His = B flat; Phe, as well as SeC = B; Arg, Tyr = sharp C; Trp = sharp D which are deduced from the notes of the code by taking the notes of the chromatic tempered scale which are

symmetrical to those of said keynotes with respect to central G.

There is another code for inhibition, which is deduced from the preceding code by symmetrization of the logarithms of the frequencies around their central value:

Trp = C; Arg, Tyr = D; Phe, SeC = E flat;  
His = E; Gln Lys, Glu Met = F; Leu, Ile,  
Asn, Asp = G; Pro, Val, Thr, Cys = A; Ser =  
B flat; Ala = sharp D; Gly = sharp F

that are deduced from the notes of the code by taking the notes of the chromatic tempered scale which are symmetrical to those of said keynotes with respect to central G.

The application of the universal code results in scaling waves respectively in phase with and in phase opposition to those taking place during the synthesis process. The term "universal code" means that this code is identical for all proteins to within the approximation of the tempered scale; the low A, for a central frequency located 76 octaves below the centre of gravity of the initial frequencies of leucine, isoleucine, and asparagine, is at 220 Hz. The expression of harmonic distance given above extends the definition suggested by Y. Hellegouarch in C. R. Math. Rep. Acad. Sci. Canada, Volume 4, Page 227, 1982. The exact values of the frequencies depend on the proportions of the groups of the above-mentioned amino acids among the transfer RNA population surrounding the protein biosynthesis.

Determination of Frequency. The next step is to derive the frequency of each of the notes. The following code is derived in the following manner, which also



optionally enables to give a more precise frequency value to each note. The frequency of the musical notes is calculated from the frequencies of amino acids in their free state (proportional to their masses) by minimizing the global harmonic distance  $\sum_{i,j} P_i P_j \log \sup (p_i, q_j)$  calculated for all possible pairs of notes,  $(p_i/q_j)$  being the harmonic intervals globally the closest to the corresponding proper frequency ratios. Their respective proportions  $P_i, P_j$  in the environing population of transfer RNAs are taken into account. While respecting the condition  $\delta f < \Delta f/2$  where  $\delta f$  is the displacement of the initial frequency towards its synchronized value and  $\Delta f$  is the interval between the two successive synchronized frequencies of the obtained scale, which encompass this initial frequency. The resulting frequency is then transposed into the field of audible frequencies. See, method described in the French patent number 8302122.

Determination Of The Musical Period. Once the frequency of each musical note is determined, the musical period is determined by identifying similar series of musical notes. The existence of musical periods results directly from that of scaling waves.

An indication is given by the presence of obvious cadences producing punctuations in the musical development. Obvious cadences include such cadences as GG, F-S. That is to say, F closely followed by S, as well as the cadence ending the signal peptide when it is present, for stimulation; series of R or Y, for inhibition; exceptionally, relative pauses induced by

harmonic variations which would otherwise be too straight; and in all cases, cadences expressing the return to the tonic note.

5 The similar passages are then determined. One method of determination is by the direct repetition of notes. When this method is used the period is given by a simple calculation of autocorrelations of notes. More specifically, by minimizing the frequency differences between notes by the number that minimizes the average on  
10 the protein of melodic distances between notes located an integer number of intervals apart.

A second method is to determine the melodic movements of the musical notes. The period is calculated by autocorrelations of signatures -- or frequency  
15 variation signs - from one note to the next. More specifically, the period is determined by calculating autocorrelations of the melodic distances from one note to the other, the distances being counted with their sign, i.e., multiplied by the corresponding signatures;  
20 or even more finely, by the number which minimizes the average on the protein of step by step melodic distances variations, to within an integer number of intervals apart. The repetition of the melodic contours are processed by a calculation of autocorrelations of pairs,  
25 or even better, of triplets of signatures.

A third method of determining the period of the musical notes is by the logic of the harmonic movement that reproduces the notes or the melodic movement to the nearest simple harmonic transposition. The period is  
30 then given by the number that minimizes the average on

the protein of harmonic distances between notes located an integer number of intervals apart.

Sometimes when an "alignment" of similar sequences is present, the period appears in the additions or in the deletions of certain of the sequences. The result gives a melodically and harmonically coherent progression. To do that, account is taken of the fact that the last notes of each period or member of phrase -- usually the second half, and more particularly the last note -- as well as those situated on the strong beat are the most important for this progression. The final result is the most significant respecting the whole of these criteria. These different elements are balanced according to their relative importance in the protein, and especially the harmonic and melodic distance by the square of the ratio of their normalized standard deviations. There is usually one that is distinctly more significant than the others.

Cases similar to allosteria nevertheless exist, and have a biological meaning (stimulation or inhibition by such molecule or such other one during the metabolism), but influence more frequently the position of the measure bars than the period. It is noted that metabolic function is different according to the context, for instance, CG rich or AT rich; the measure bars depending upon the composition of the DNA, as the "Christmas trees" that can be seen during certain syntheses clearly displayed (cf. B. Alberts and al., *Molecular biology of the cell*, 2<sup>nd</sup> edition, Garland Publ. Co. 1989, page 539).

Determining The Lengths Of Musical Notes. If necessary, the period is rectified so that the melodic

passages that repeat or follow one another can be found in the same place inside the measure. From this rectification the individual lengths of the musical notes are deduced. This operation of adjusting the phrasing to the measure is comparable to the well known phenomenon of lengthening the vowels of a sung text.

In practice, the operations described above can be performed most easily with a keyboard, such as a Casio™ equipped with a "one key play" device, or with a computer programmed especially for that purpose with stored sequence of musical notes and where the sequence of notes can be played. However, some precautions are required. Prudence implies, among other things, to decode the same molecule or a musically similar molecule, in the direction of inhibition (or in any case in the direction opposite from the initial one), taking into account the fact that molecules very often have a preferential decoding direction. It is often the case that pairs of molecules that sensibly exert the same function find one pair being more musical in inhibition and the other one in stimulation.

Checking Rhythmic Style Through The Distribution Of The Bases of DNA. When the molecule is musical enough, the period of autocorrelations corresponds to that of the protein. The autocorrelations determine in principle the measure bars, the ranks of base triplets -- or more precisely of bases in third position in these triplets -- for which the peaks of autocorrelation are the highest, corresponding to the most accentuated notes. By referring to codon sage, in comparison with known molecules (already decoded, or more regular and thus

raising less difficulties) having the same supposed  
rhythmic style; the style of musical rhythm (which by  
constraining the accentuation of notes, influences the  
choice of bases in third position) determining the codon  
usage. Molecules of the same style must therefore have  
the same codon usage. If necessary, the decoding of some  
passages is corrected.

Determining The Tone Quality. Tone quality is, in  
principle, different for every molecule and for every  
distribution of musical notes. In theory, tone quality  
mainly depends upon the molecule itself but it also  
depends upon all the levels of the organism which  
retroact on the harmonic structure of amino acid  
vibrations. The tone quality of the musical sequence is  
determined by comparing the repartition of the music  
sequence of the amino acid chain to the average  
repartition of those notes of the whole of the protein to  
determine which harmonics must be raised or lowered. The  
term "tone quality" or timbre is characterized by the  
harmonic structure of a note and more precisely by the  
variation of harmonic structure over a given note.

A first approach is given by adjusting the  
distribution of molecule notes to the theoretical graph  
of that distribution. The distribution is deduced from  
the scaling wave equation. The distribution also  
corresponds to what can be observed in average, on the  
whole of proteins. This adjustment to the tone quality  
requires determination of which harmonics are amplified  
and which are softened in the wanted tone. See, French  
Patent No. 8302122. The closest tone quality is then  
selected in a palette of given ones. For example, a voice

memory or as one can already find included in many  
expanders and musical softwares. To distinguish more  
precisely between three situations: (1) distribution of  
notes constant along the molecule to provide a relatively  
5 fixed harmonic structure; (2) straight distribution  
changes to provide different successive tones of  
instrument, for instance cytochrome C with several organ  
registers; and (3) progressive distribution change which  
then reproduces the time evolution of the harmonic  
10 structure of one note, for example, myosin, where this  
evolution indicates a timbre of trumpet.

Apart from this, determining the tempo gives no real  
problem to the technician because it normally follows  
from the rhythmic style. It is generally all the faster  
15 that there are important redundancies in the proteic  
sequence, as it is the case for fibrous proteins.

Determining The Colors. Optionally, the colors are  
determined by applying the universal code. The color is  
deduced from vibration frequencies of individual amino  
20 acids through the formula (drawn from scaling wave  
theory):  $\nu \sim \nu_0 \operatorname{Argch} (e (f/f_0) \operatorname{Logch} 1)$ , where  $(f, f_0)$   
represent the proper quantum frequencies associated with  
aminoacids as previously, and  $\nu, \nu_0$  those of colors, the  
index  $_0$  showing central values. This gives the following  
25 code relating to the stabilization of proteins  
synthesized *in situ* (the code related to the  
stabilization of their inhibition is deduced as in  
section 1 by symmetrization of the logarithms of  
frequencies with respect to the central lemon yellow):

Gly = dark red: Ala = bright red: Ser =  
orange; Pro, Val, Thr, Cys = ochre; Leu,  
Ile, Asn, Asp = lemon yellow; Gln, Glu, Lys,  
Met = green; His = emerald: Phe = blue; Arg,  
5 Tyr = indigo; Trp = purple,

these frequencies then being moved towards red or purple  
according to the global repartition of the molecule  
frequencies in a way similar to the description for tone  
quality as above. The spatial position of colors is the  
10 same as those of the amino acids in the tridimensional  
spatial representation of the molecules.

Several examples are set forth below to  
illustrate the invention and the manner in which it is  
carried out. In these examples as well as in the  
15 figures, the one-letter notation for amino acids:  
Gly = G; Ala = A; Ser = S; Pro, Val, Thr, Cys = P, V, T,  
C respectively; Leu, Ile, Asn, Asp = L, I, N, D; Gln,  
Glu, Lys, Met = Q, E, K, M; His = H; Phe = F; Arg, Tyr =  
R, Y; Trp = W is used.

#### 20 **Example 1**

This example illustrates decoding a protein that is  
regular from beginning to end. Cytochrome C provides a  
constant deletion of eight amino acids (sometimes seven)  
among animal proteins when compared to plants. Observing  
25 the autocorrelations of musical notes and melodic  
contours confirmed the value of the musical period.

The occurrences of the same note was counted and the  
same direction of pitch variation occurred three times in  
a row (the same triplet of signatures), which was distant

from an integer number k of musical notes. The following result was obtained:

Values of k	1	2	3	4	5	6	7	8	9	10	11	12
Note autocorrelations	19	15	15	20	19	15	17	21	14	17	18	13
5 Melodic contour autocorr.	1	7	4	6	5	10	8	13	5	4	4	4
Total	20	22	19	26	24	25	25	34	9	21	22	17

the peak at k = 8 being worth about 2.5 standard deviations (as compared to its expectation value  $22.3 \pm 4.7$  determined from the repartition of notes of the molecule). The significance of this peak was reinforced when using melodic distances.

The peak outgrew distinctly 3 standard deviations when the autocorrelations of melodic intervals were included by taking as a definition of the melodic distance between two notes, the absolute value of the difference of the ordinal ranks of their tempered frequencies arranged in ascending order. This definition is derived from the usual nomenclature: second, third, etc., for the notes of a musical mode. The secondary peak at k = 7 then became slightly significant, corresponding to the relative stretching of the seventh note which tended to precede the return to the tonic; whereas, the one at k = 4 was reinforced when harmonic distances were used to spatial foldings of the molecule.

The observation of the cadences also confirmed this value, as well as that of the internal similarities. The last five notes of the first, second and third group of eight produced together an exact harmonic superposition. In other words, a canon for three voices. More precisely, these last two investigations showed a greater



relative importance of the seventh note (F-S cadence on  
the second period) and the eighth note (back to the A  
minor tonic) for each period. The latter once more  
prevailing over the former. That is, the perfect S-Q  
5 cadence on the sixteenth note prevailed over the  
preceding F-S cadence with the recovering of the initial  
tonality. The division of the period resulted in six  
semiquavers, one quaver, one crotchet (which meant  
relative lengths 1-1-1-1-1-1-2-4 with a 6:8 rhythm as  
10 shown in Fig. 1). The coherence of the melodic  
progression (wherefrom the observed regularity mainly  
proceeds) as well as the richness of the harmonic  
progression, the A minor tonality being accompanied with  
modulations in E minor (second bar), G minor (eighth  
15 bar), and F major (third and ninth bar) was apparent.

The first and seventh notes of each period fostered,  
respectively, adenine and thymine in third position;  
whereas, the third and eighth notes fostered in the same  
way cytosine and guanine. This confirmed the above  
20 division for the period and the relative lengths of  
notes. In other words, the seventh and eighth notes had  
lengths that were respectively twice and four times the  
first. This also showed that in an AT-rich environment  
strong beats were on the first and seventh notes, and  
25 therefore the measure bars were on the first. However,  
in a CG-rich environment the musical sequence started on  
an anacrouse (strong beat on the third and eighth notes,  
measure bar on the third).

The conclusion was that the protein had distinct  
30 metabolic roles, depending on its environment.

Actually, the range of its metabolic action was first demonstrated by the degree of its musical evolution. In comparison with the sequence of *Euglena gracilis*, in the three first measures an improvement of 56% of the melodic [regularity] level and of 16% of the harmonic [regularity] level was observed as defined from the minimization of the respectively melodic and harmonic distances between successive notes.

The search of musical similarities with other proteins showed the possibility to superpose cytochrome C onto endozepine with a musical reading frame compatible with the measure bar on the first note. This resulted in a slightly AT-rich molecule; thereby predicting an anti-depressive role for the cytochrome (and its music), through the eventual desinhibition of neurotransmission; as well as, a musical enchainment (then beginning on an anacrouse) with cytochrome oxidase. Cytochrome oxidase is slightly CG-rich and ends the respiratory chain.

As for tone quality, because tonality was present in A (minor), the quasi absence of the fourth (D) and the relative weakness of the fifth (E) compared to the distinct dominance of the tonic note and to the abundance of the octave (low A - medium A) privileged harmonics 1 and 2, to the prejudice of the followings, indicated an organ timbre with slightly different registers according to the passages.

As shown in Fig. 2, colors effectively grouped themselves into colored stains onto the mature protein with, as in the case for music, remarkable harmonic responses. The color determination was useful to confirm the musical decoding, insofar as some autocorrelations of

notes were translated not into the musical period but in the spatial folding of the molecule. The spatial folding must eventually be subtracted to determine the musical periods. It was found that where a secondary peak of these autocorrelations,  $k = 4$ , due to the  $\alpha$ -helix of the beginning which can be seen in Fig. 2, corresponded to these foldings. Conversely, the musical decoding gave indications about the spatial structure of a protein.

### Example 2

This example illustrates control of the decoding of a protein showing rhythmical variations. The decoding was controlled at different levels including the decoding of molecules known to be metabolically agonist and the coherence of the conclusions that were drawn from the musical similarities observed.

Recovering full sections of the metabolism facilitates the decoding. In Example 1, the "rhythmic formula" of cytochrome C was transcribed as follows:

|6/8 GDVEKGK:K::|IFIMKCS:Q::|CHTVEKG:G::|, etc.

+ + + + + + + + + + + + + + + +

where the + underline the strong beats, the | indicate the place of measure bars and the : indicate the lengthening of notes.

In subunit III of cytochrome oxidase, which is musically chained to cytochrome C, the beginning is a four-time formula as shown by the internal similarities. The notes 7 to 22, which remind in their contours the manner of Bach, were split into groups of four notes, each one being superposable to the next. At the tenth measure, another measure which was not only superposable

was found onto the first measure of cytochrome C, but was  
in fact, even practically identical to the third measure  
of the same cytochrome. This implied a lengthening of the  
eighth measure (as the cadence seen at the end of this  
5 measure already indicated in itself), in a six-time  
measure (Fig. 1):

```
      | 4/8 MTHQSHAY | HMKPSPW | PLTGALSA | LLMTSGLA |  
      + + + + + + + + + + + + + + + + + + + + + +  
      MWFHFHSM | TLLMLGLL | TNTLTMYQ || 6/8 WWRDVTR: : : : : |  
10    + + + + + + + + + + + + + + + + + + + + + +  
      ESTYQGH: H: : : : | TPPVQKG: : : : : ||  
      + + + + + + + + + + + + + + + + + + + + + +
```

This change in rhythm (from 4/8 to 6/8) was visible in  
base autocorrelations of the DNA where, at this point,  
15 the prominent peak went from the fourth to the sixth base  
triplet.

As seen in Fig. 1, the sequence started on an  
anacrouse emphasizing the strong beat on the third note,  
in view of the enchainment with the CG-rich rhythmic  
20 variant of cytochrome C.

### Example 3

The example illustrates reconstitution of a  
metabolic chain including stimulations and inhibitions.

The decoding of histone 4 was particularly easy.  
25 The periodicity of 7 is clearly visible on the sequence  
at the outset of the molecule. The repetition of G  
within a two amino acid interval indicates a binary  
rhythm, and the GG cadences that end the two first  
periods specify right away a four-time rhythm:

```
30    | SGRGKGG: | KGLGKGG: | ;
```

+ + + + + + + +

this pattern continued until the end of the sequence,  
with the only exception being the last measure which was  
syncopated to recover the rhythm of the first two  
5 measures. See Fig. 3. The global repartition of the  
notes showed a harmonic structure corresponding to the  
tone of a flute. The "skip of notes" repeated from the  
beginning, which suggested a sound with an attack and a  
timbre similar to that of Pan's pipes.

10 Histone 4 is one of the most conserved proteins  
among the animal and plant kingdoms. This does not mean  
that its metabolic action doesn't sometimes need to be  
tempered. The theme of histone 4's first two measures  
was found in inhibition and transposed to the fourth, in  
15 the conserved part of the beginning of chalcone synthase,  
which is the pigmentation enzyme of many flowering  
plants. See Fig. 3. This may be compared to the supposed  
role of chromatin, which histone 4 is part of, in the  
process of magnesium fixation. During spring, plants  
20 need a lot of magnesium for photosynthesis and the  
plant's fixation needs to be stimulated. Chalcone  
synthase is then inhibited; whereas, during the fall, the  
weaker stimulation of histone desinhibits chalcone  
synthase and allows the replacement of the green of the  
25 leaves by brighter colors of that season, the diversity  
of which, so much praised by the poets, becomes thus more  
understandable through their epigenetic component.

When listening to the musical transposition of  
histone 4, several auditors reported "an urge to eat  
30 chocolate" which contains magnesium. Some auditors found  
that "it produces the same effect as that of granulated

66000-444444  
magnesium, except that this effect is immediate in this case". This presents some inconvenience for people having a slightly too high rate of cholesterol.

Actually, the musical decoding of chalcone isomerase -  
5 the metabolically agonist of chalcone synthase, but which "works better" musically in stimulation -- included a series of themes and variations whose succession reproduced, in flowering plants, themes of the full metabolic chain regulating cholesterol in man. In  
10 addition, the frequency of the ascending fourths in chalcone isomerase tended to approximate that observed in the alkali light chain of mammalian myosin, which stimulated muscular contraction (while magnesium acted as a muscular decontractant). Listening to the musical  
15 transposition of histone 4 encouraged physical exercise which is another way to lower cholesterol.

In fact, this example underlines the importance of a quasi-general phenomenon, that is, the epigenetic co-operation of different factors in the stimulation of  
20 protein synthesis, which accounts for the aspect meaningful in itself of the musical sequences. In this way for example, listening to myosin will generally suggest a military march.

#### Example 4

25 This example illustrates the biochemical analysis of an epigenetic cooperation involving harmonic superpositions. The biochemical analysis of these epigenetic cooperations is a valuable help for decoding.

Another way to stimulate epigenetically the muscular  
30 decontraction is heat, whose healing action for

reumatism, for example, is well known. The action of heat is conveyed by a group of proteins called heat shock, generally synthesized together. This suggests that the proteins should show harmonic superpositions.

5 In fact, the hsp 27 protein, which appeared to be the most musical, superposed itself onto the beginning of the hsp 70 protein, the most abundant, which sort of played here the role of a bass line. These two molecules were again superposable together with the beginning of  
10 troponin C, which regulates calcium in muscular contraction. The conclusion was that it plays a role as an anti-rheumatic and that its musical level is high (Fig. 4). Other molecules, also of a high musical level and epigenetically sensitive, were implicated in this  
15 type of ailment, from the stimulation of prolactin and beta-lipotropin (precursor of beta-endorphin) to the inhibition of estrogen receptor, including the inhibition of IgE and interleukin 1 beta.

These examples clearly show how large sections of  
20 the metabolism can be reconstituted step by step, with many ways to check or control the coherence of the results obtained, and thereby to precise the musical decoding of the concerned proteins.

#### 25 **Example 5**

This example shows a practical application of the method of this invention using the transcriptions in the form of either musical scores, or of recordings of the obtained musical sequences.

5 The recordings of musical sequences may be realized from musical scores described earlier, by using one of the methods evaluated in B. H. Repp, J. Acoust. Soc. Am. 88, p.622 (1990). The most precise of these methods was used in the examples hereby given.

10 In the fields of agronomy and textile industries this invention provides methods to stimulate certain specific protein synthesis, for example, bovine lactation, fermenting of baker's yeast, the sweet taste of some fruits, animal or plant fibres (keratine of sheep's wool, fibroin of silkworm, etc.), as well as the proteins specific to certain medicinal plants. In the field of environment the method of this invention is used, for example, in the assimilation of industrial effluents through plants by stimulating the biosynthesis of the corresponding proteins.

15 The method of this invention was used on a cow who regularly, during 15 days and at the time of milking, listened to recordings of musical transcriptions of the amino acid sequences of bovine prolactin, lactoglobulin, and lactalbumin. A reduction, by a ratio of 3, of the relative quantity of whey was observed, resulting in a milk highly enriched in proteins, and in a particularly savory cheese.

25 In another experiment growing tomato plants were given a "cocktail" of musical transpositions of different proteins including: specific virus inhibitors, various extensions, then a flowering enzyme (LAT 52), an antibacterial protein having musical similarity to 30 thaumatin, an improvement of sugar percentage (P 23), and inhibitors of fruit softening enzymes (pectinesterase and



polygalacturonase). A distinct increase in size and number of fruits (summing up to a ratio of about 3.5) was observed, as well as, a sensitive increase of the sweet taste in a significant proportion of the fruits that had particularly received P 23.

These noteworthy results go along with a certain amount of precautions, namely, there exist some counter-indications to an excess of stimulation, especially of prolactin, which must be cautiously taken into consideration by breeders that carry out these methods, as well as for the animals themselves who may be fragilized. In the experiments carried out on cows with Mozart music -- bovine prolactin has in fact, apart from a "musical level" particularly high which can here define in a mathematically simple way some musical turns that can be qualified as "typically Mozartian" -- the rate of mammites could seem worrying. In such a case one ought to complete the hearing of prolactin with that of alpha-1 antitrypsin, whose musicality is also very elaborate and whose metabolism is complementary. Similarly for tomatoes receiving outside stimulations, one must be cautious not to interrupt the cycle too suddenly.

These results give an indication of the order of magnitude of results obtainable in such conditions.

#### **Example 6**

In the therapeutic and preventive fields, many ailments are characterized by a specific metabolic weakness and can therefore be efficiently prevented or treated with the help of the present invention. This example illustrates such prevention or treatment.

Because the minimal length of a musically active sequence is of the order of that of a signal peptide, i.e., from several amino acids to a few tens, this action may be very fast and appear after a few seconds or a few minutes. Nevertheless, the complete integration of the produced effect can take slightly more time, or even require, in case of a strong cultural conditioning, i.e., a certain initial training. But usually, this initial training is accomplished rather rapidly for the obvious benefit of the persons concerned.

For a responsible use of the described method, it is important to know the metabolic role of the molecules involved. And it is of course one of the interests of the musical decoding of proteins (associated to the corresponding colors) to allow, by systematically spotting the similarities and counter-similarities of melodies (and colors) from the protein sequences that are known and available in data banks, to select proteins that are metabolically agonist and antagonist of a given protein, for which the degree of musical elaboration also gives an indication of the importance of its metabolic role. The described method therefore allows determinations of precise particular indications for some proteic sequences.

As earlier noted, in animal or plant proteins, especially among the most musical ones, successive melodic fragments of human metabolic chains were observed. Therefore, the transpositions which were found to be active on man were not limited to human molecules. On the other hand, the metabolism of those species seems in some way more "specialized" for the

production of certain molecules, and it is indeed the  
most musical proteins that will be the most important for  
the applications. Of course, these correspondences  
between different species facilitate the delimitation of  
5 the metabolic role, and the decoding of proteic  
sequences.

The musicality of a molecule implies in itself that  
its epigenetic stimulation is preferable for a  
therapeutic use, (because of the range of its metabolic  
10 interactions), to its direct absorption. The "most  
musical" molecules are generally those for which either  
the production by genetic engineering, or the therapeutic  
use which derives from it, will meet some problems, such  
as of transportation to the site of action, or of  
15 stability, or more specifically of secondary effects  
related to doses that should be much more important than  
what they are in the body to obtain comparable effects,  
because then, the scaling waves naturally associated to  
their production are not present any more. This is  
20 particularly true for the inhibition of proteins, when  
the natural inhibitor is much heavier, or simply when the  
production needs to be reduced at a given time or in a  
systematic way.

Eventually, concerning the use of transcriptions of  
25 proteic sequences, the very quickness of their action may  
allow, by differential comparison, especially bipolar, of  
their positive and negative effects to precisely which  
one is the most appropriate in a given situation. This  
identification is facilitated by the comparison with  
30 transcriptions of known proteic sequences of acoustic or  
electromagnetic phenomena exhibiting distinct series of

frequencies, and for which some effects have been observed in a similar situation.

5 As will be appreciated from the above, the invention is in no way limited to those methods of putting it into effect, of construction and of application which have been described above in detail; on the contrary, it covers all versions which may be conceived of by workers skilled in the art, without exceeding, either the framework or the scope of the present invention.

THAT WHICH IS CLAIMED IS:

1. A method of regulating protein synthesis *in situ* comprising:

5 (a) determining the sequence of musical notes associated with the amino acid chain of a protein by associating with each amino acid a musical note whose frequency is transposed from the proper frequency of the amino acid;

10 (b) determining the musical periods of said sequence of musical notes by identifying similar series of musical notes;

15 (c) comparing the repartition of said musical sequence of said amino acid chain to the average repartition of said musical notes of the whole of proteins so as to determine the tone quality; and

(d) regulating the biosynthesis of said protein by playing said sequence of musical notes, including the musical period of said notes and the tone quality of said musical notes.

20 2. The method of regulating protein synthesis according to Claim 1 further comprising:

25 determining the lengths of said musical notes by rectifying collectively, and then rectifying individually said musical periods by adjusting the phrasing to the measure of said musical sequence.

3. The method of regulating protein synthesis according to Claim 1 further comprising determining the frequency of said musical note according to a code comprising:

(a) taking the frequency of each amino acid in its free state, proportional to its mass,

(b) minimizing the global harmonic distance between the frequencies of each pair of amino acids in said protein while taking into account the proportion of each amino acid in the population of transfer RNAs within a cell where synthesis of said protein takes place, and wherein the displacement of the note frequency towards its synchronized value is inferior to half the interval between the two synchronized frequencies which surround said keynote frequency, then

(c) transposing the frequencies thus obtained into the auditive range, said code being relative to the biosynthetic stimulation of said protein; and

(d) obtaining said code relative to its inhibition by symmetrization of the logarithms of heretofore obtained frequencies with respect to their central value considered as the origin.

4. The method of Claim 3, wherein said code comprises the following notes of the chromatic tempered scale in ascending order:

Gly = low A; Ala = C; Ser = E; Pro, Val, Thr, Cys = F; Leu, Ile, Asn, Asp = G; Gln, Lys, Glu, Met = A; His = B flat; Phe as well as SeC = B; Arg, tyr = sharp C; Trp = sharp D.

5. The method of Claim 3, wherein said code comprises the following notes of a chromatic tempered scale, in ascending order:

Trp = C; Arg, Tyr = D; Phe as well as SeC =  
E flat; His = E; Gln, Lys, Glu, Met = F;  
Leu, Ile, Asn, Asp = G; Pro, Val, Thr, Cys =  
A; Ser = B flat; Ala = Sharp D; Gly = sharp  
F,

which are deduced from the notes of the code by  
taking the notes of the chromatic tempered scale which  
are symmetrical to those of said keynotes with respect to  
central G.

6. The method according to Claim 3 wherein said  
synthesis stimulates synthesis of a protein in a plant.

7. The method according to Claim 1, wherein each  
sound transposition of quantum vibrations associated with  
the biosynthesis of a given protein is completed by the  
color transposition of quantum vibrations associated to  
the mature protein after it is spatially folded back over  
itself, according to a code specific to the stabilization  
of that protein or to the inhibition of its biosynthesis  
obtained through the musical sequence realized according  
to Claim 1, which code is deduced from the code obtained  
from Claim 1, by application of the formula  $\nu \cong \nu_0$   
 $\text{Arch}(e^{f/f_0})^{\text{Logch } 1}$ , where  $f, f_0$  are the musical frequencies  
and  $\nu, \nu_0$  the frequencies of colors, with the index  $_0$   
showing the central values.

8. The method according to Claim 6, wherein the  
stabilization of proteins stimulated by the musical  
sequences obtained according to Claim 1 consists in the  
association to the different amino acids of the following  
colors:

Gly = dark red; Ala = bright red; Ser = orange;  
Pro, Val, Thr, Cys = ochre; Leu, Ile, Asn, Asp =  
lemon yellow; Gln, Glu, Lys, Met = green; His =  
emerald; Phe = blue; Arg, Tyr = indigo; Trp = purple

5           9.    Transcriptions of a musical sequence according  
to Claim 1 selected from the group consisting of musical  
scores of said musical sequence and audio recordings of  
music according to said musical sequence.

10           10.   The method according to Claim 9 for the  
characterization of proteic sequences fit to be regulated  
by using any of the transcriptions characterized in that  
one delimits their metabolic role by decoding with the  
method according to Claim 3, thereby showing the musical  
similarities and anti-similarities that they present with  
15   other proteins, the harmonic superpositions with other  
proteic melodies, or a combination of these factors, from  
which the agonisms and antagonisms can be deduced.

20           11.   The method according to Claim 9, in which the  
characterization for a given application is refined by  
bipolar differential comparisons with the positive or  
negative effects obtained by using said transcriptions.

25           12.   The method according to Claim 10, in which the  
characterization for a given application is refined by  
identification, through musical similarity or anti-  
similarity of the proteins involved during positive or  
negative effects due or associated to acoustic or  
electromagnetic phenomena exhibiting distinct series of  
frequencies.



METHOD FOR THE REGULATION OF PROTEIN  
BIOSYNTHESIS

ABSTRACT OF THE DISCLOSURE

5           There is provided a method for determining the  
musical notes associated with an amino acid sequence, the  
musical periods of the sequence, the lengths of the  
notes, and the tone quality of the notes through the  
retroaction of the whole set of amino acids and using  
10   that information to regulate the biosynthesis of the  
protein. The amino acids that build a protein emit a  
signal of quantum nature at a certain frequency.  
Following the properties of this signal, the frequency is  
transposed into a musical note. This discovery has  
15   numerous applications since one can then deduce from the  
amino acid sequence of a protein a sequence of notes  
composing the melody that will act to stimulate or  
inhibit its synthesis inside an organism, wherefrom one  
can in addition delimit its biological functions.

20  
CLT01/4353653v1

# Cytochrome oxidase

Last protein of human  
respiratory chain

Beginning of mitochondrial  
subunit 3



# Cytochrome C


One before last protein  
of human respiratory chain

Complete sequence



Figure 1

669250-26902660



## Symbols used

|   |   |   |   |   |   |   |   |   |   |   |
|---|---|---|---|---|---|---|---|---|---|---|
| ○ | ⊗ | ⊙ | ⊗ | ⊙ | ⊗ | ⊙ | ⊗ | ⊙ | ⊗ | ⊙ |
| G | A | S | P | L | B | H | F | R | W |   |
|   |   |   | V | I | K |   |   |   | Y |   |
|   |   |   | T | N | E |   |   |   |   |   |
|   |   |   | C | D | M |   |   |   |   |   |

Human cytochrome C  
Amino end region

Figure 2

## Histone IV

Protein of human chromatin

Complete sequence



## Chalcone synthase

Pigmentation enzyme  
of Petunia hybrida flowers

IN INHIBITION  
Beginning



Figure 3

## 'Heat shock' hsp 27 and hsp 70

Human proteins  
synthesized under the effect of heat

Beginning

M T E R R V P F S L L R G P S W  
M A K A A A V G I D L G T T Y S

## Troponin C

Regulates calcium  
in human skeletal muscle

Complete sequence

DTQQAARSYLSEEMIAEFKAAFD MFDA DGGGDIS  
VKELGTVM RMLGQTPTKEELD AII EEVD E DGS  
GTIDFEEFLVMMVRQM KEDAKGKS EEELAE CF  
RIFDRNADGYIDPEEL AEIFRASGEHVTD EEI  
ESLMKDGDKNNDGRIDFDEF LKMM EGVQ

Figure 4

## DECLARATION AND POWER OF ATTORNEY FOR PATENT APPLICATION

Attorney Docket No. 3339-239A

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled

### METHOD FOR THE REGULATION OF PROTEIN BIOSYNTHESIS

the specification of which

☒ is attached hereto

OR

☐ was filed on \_\_\_\_\_ as United States Application No. or PCT International Application Number \_\_\_\_\_ and was amended on \_\_\_\_\_ (if applicable).

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is material to patentability as defined in Title 37 Code of Federal Regulations, § 1.56.

I hereby claim foreign priority benefits under Title 35, United States Code, § 119(a)-(d) or § 365(b) of any foreign application(s) for patent or inventor's certificate, or § 365(a) of any PCT international application which designated at least one country other than the United States of America, listed below and have also identified below any foreign application for patent or inventor's certificate, or of any PCT International application having a filing date before that of the application on which priority is claimed.

|          |         |                  |   |
|----------|---------|------------------|---|
| 92 06765 | France  | June 4, 1992     | <input checked="" type="checkbox"/> Yes <input type="checkbox"/> No |
| Number   | Country | MM/DD/YYYY Filed | Priority Claimed  |
|          |         |                  | <input type="checkbox"/> Yes <input type="checkbox"/> No            |
| Number   | Country | MM/DD/YYYY Filed | Priority Claimed  |
|          |         |                  | <input type="checkbox"/> Yes <input type="checkbox"/> No            |
| Number   | Country | MM/DD/YYYY Filed | Priority Claimed  |

I hereby claim the benefit under Title 35, United States Code, § 119(e) of any United States provisional application(s) listed below.

|                    |                          |
|--------------------|--------------------------|
|                    |                          |
| Application Number | Filing Date (MM/DD/YYYY) |
|                    |                          |
| Application Number | Filing Date (MM/DD/YYYY) |

I hereby claim the benefit under Title 35, United States Code, § 120 of any United States application(s) or § 365(c) of any PCT international application designating the United States of America, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT International application in the manner provided by the first paragraph of Title 35, United States Code, § 112, I acknowledge the duty to disclose information which is material to patentability as defined in Title 37, Code of Federal Regulations, § 1.56 which became available between the filing date of the prior application and the national or PCT international filing date of this application (37 C.F.R. § 1.63(d)).

|                 |                  |                                      |
|-----------------|------------------|--------------------------------------|
| PCT/FR 92/00524 | June 2, 1993     | National Phase Entered               |
| Application No. | Filing Date      | Status<br>Patented/Pending/Abandoned |
| 08/347,353      | December 1, 1994 | Pending                              |
| Application No. | Filing Date      | Status<br>Patented/Pending/Abandoned |
|                 |                  |                                      |
| Application No. | Filing Date      | Status<br>Patented/Pending/Abandoned |

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

POWER OF ATTORNEY: As a named inventor, I hereby appoint the practitioners associated with the Customer Number provided below to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith, and direct that all correspondence be addressed to that Customer Number:

**Customer Number 000826**

Direct correspondence to the  
attention of and telephone calls to:

Ernest B. Lipscomb, III  
Registration No. 24,733  
Tel Charlotte Office (704) 331-6000  
Fax Charlotte Office (704) 334-2014

Full name of (first/sole) inventor: Joel Sternheimer

Inventor's

Signature: Joel Sternheimer JSH Date: 5/19/99

Residence:

Citizenship: French

Post Office Address:

CLT01/4358048v1

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re: Joël Sternheimer  
Appl. No.: To Be Assigned  
Filed: Concurrently Herewith  
For: METHOD FOR THE REGULATION OF  
PROTEIN BIOSYNTHESIS

May 25, 1999

Assistant Commissioner for Patents  
Washington, DC 20231

**RULE 132 DECLARATION  
OF JOËL STERNHEIMER**

Sir:

I, JOËL STERNHEIMER, do hereby declare and say as follows:

1. That I am a graduate of Paris, Lyons and Princeton Universities and received my degrees in the years 1964, 1966 and 1967 (Doctorate in 1966).  
  
My Curriculum Vitae and the list of publications are attached thereto (Annexes 1 and 2).
2. That it clearly emerges from the Invention as now defined that it does not lack utility: the method of the invention allows and controls the *in situ* regulation of the synthesis of selected protein. It was undertaken by Mr. Pedro Ferrandiz under my supervision to stimulate the growth of blue-green algae –prokaryotes, genus *Anabaena*- by epigenetic regulation. Their photosynthetic activity involves in particular pigmentary proteins (cyanins). Thus their biosynthesis is easily observed through color change and oxygen release.



We want to point out that this first attempt of stimulation in an aquatic medium is relatively simple to reproduce. We believe that the results obtained are particularly promising. One may add the fact that it points towards numerous applications.

- Materials and methods

- Dilution of 12 ml of *Anabaena variabilis* (stock provided by the Ecole Normale Supérieure de Paris) in 1 500 ml of mineral water.
  - Addition of 40 g of dry vegetable manure containing 8%, say 2.6 g, of nitrates as well as 40 g of river pebbles (as suggested by Vincent Bargoin this would provide the solution with trace elements).
  - Adaptation time to the cultures medium: four days.
  - Transfer of 750 ml of the solution in two vats subjected to natural enlightenment.
- This setting in culture started on the 30<sup>th</sup> of April.

*Musical diffusions.*

The music has been diffused in one of the vats, by mean of an aquatic speaker Altec UW-30, while the other vat served as a control.

The proteins transcribed in musical sequences were the following:

- TAPE I (45 min)
  - NIF H of *Anabaena v.* (five times)
  - Allophycocyanin of *Anabaena v.* (three times)
  - Plastocyanin of *Anabeana v.* (three times)
  - Nitrate reductase of *Chorella s.* (three times)
  - PS1 photosystem protein of *Anabaena v.* (three times) (\*)
  - Ferredoxin of *Anabaena v.* (five times)
  - Protein 35 K of *Anabaena v.* (eight times) (\*).
- TAPE II (15 min)
  - Allophycocyanin of *Anabaena v.* (two times)
  - Plastocyanin of *Anabaena v.* (two times)
  - PS1 photosystem protein of *Anabaena v.* (three times) (\*)
  - Ferredoxin of *Anabaena v.* (four times)
  - Protein 35 K of *Anabaena v.* (eight times).
- TAPE III (15 min)
  - Ferredoxin of *Anabeana v.* (two times)
  - NIF H of *Anabeana v.* (three times)

NIF A of *Anabaena* v. (three times) (\*)

NIF D of *Anabaena* v. (three times) (\*)

Nitrate reductase of *Chlorella* s. (three times)

Protein 35 K of *Anabaena* v. (two times) (\*).

The transcriptions had been realized by J. Sternheimer on a sampler Casio SK1 apart from those labelled (\*) which were made by P. Ferrandiz on a « One Key Play » software written by Sylvie Guillou and Fabrice Ocelli (INSERM St-Anne, Paris). The rate of the transpositions is tuned so as to make their length correspond to the photoperiods of the micro-organisms.

Tape I was played twice a day, from the 30<sup>th</sup> of April to the 5<sup>th</sup> of May. Then from the 7<sup>th</sup> to the 10<sup>th</sup> of May TAPE II was played in the morning while TAPE III was played in the evening.

During this period the viability of the micro-organisms was regularly controlled: Samples were drawn from the cultures and then checked under a microscope.

- Results (Annex 3)

- Evolution of the coloration of cultures (Figure 1)

One poured in the vats the solutions looked opaque (after tossing).

This was due to the manure mentioned above, the dilution rate of the original stock but also to the spread of a fibrous contaminant which was not characterized.

From the 2<sup>nd</sup> day of listening the musical vat presented a greater proportion of suspending matter than the control one. However this trend reversed itself by the 4<sup>th</sup> day. We therefore assumed that the musical exposures had been too long and we decided to abort the diffusion of Tape I.

Instead Tapes II and III have been used. We then observed on May 8<sup>th</sup> that the tint of the cultures in the musical vat displayed a green blue coloration more pronounced than those in the control vat (Figure 2).

This trend kept increasing up to the end of the experiment.

- Oxygen release.

Ten days after the end of the period of diffusion the musical cultures became characterized by a proliferation of bubbles at the surface (Figure 3).

Since these bubbles had the property to revive the flame of a lighted match which was put close by, we concluded they contained oxygen. On May 24<sup>th</sup> there were about 70 surface bubbles and on the 28<sup>th</sup> they were 130 (Figure 4). We point out that the maximum number of visible bubbles observed in the control vat is 8. Hence there is more than a factor 16 between the two cultures with respect to oxygen release. In fact the medium of the musical culture was saturated with oxygen at the end of the observation time. Clearly this is correlated to an increase of the photosynthetic activity in the musical vat. It indicates that while the oxygen was released some carbonated composites have been fixed (Figure 5, taken six months later). Thus this particular application of the epigenetic regulation process led to an interesting depollutive system. This should beget further interests.

Other experiences showing the utility of the instant invention are herewith attached (Annexes 4-8).

As regards a garden experience: See Annex 9.

Figure 1 of this Annex 9 also attached is a comparative test:

On the left side (control): non treated tomatoes

On the right side: tomatoes having received during 16 days 3 minutes per day, the music of protein of anti-drought protein TAS14. Both control tomatoes and the treated tomatoes having 1 1½ litres of water per plant per day.

3. The undersigned petitioner declares further that all statements made herein of his own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of this application or any patent issuing thereon.

4. Further declarant saith not.

Joël Sternheimer JSN

Joël STERNHEIMER

1999/5/19

Date

2025-05-19 14:00:00